

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/55, 31/495, C07D 401/14, 403/14	A1	(11) International Publication Number: WO 97/17973 (43) International Publication Date: 22 May 1997 (22.05.97)
(21) International Application Number: PCT/US96/18342 (22) International Filing Date: 12 November 1996 (12.11.96) (30) Priority Data: 60/006,572 13 November 1995 (13.11.95) US (71) Applicant (for all designated States except US): SMITHKLINE BEECHAM CORPORATION [US/US]; Corporate Intellectual Property, UW2220, 709 Swedeland Road, P.O. Box 1539, King of Prussia, PA 19406-0939 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): BHATNAGAR, Pradip, Kumar [US/US]; 300 South Balderston Drive, Exton, PA 19341 (US). HEERDING, Dirk, Andries [NL/US]; 1 Mill Creek Lane, Malvern, PA 19355 (US). (74) Agents: HALL, Linda, E. et al.; SmithKline Beecham Corporation, Corporate Intellectual Property, UW2220, 709 Swedeland Road, P.O. Box 1539, King of Prussia, PA 19406-0939 (US).		(81) Designated States: JP, US, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>
(54) Title: HEMOREGULATORY COMPOUNDS (57) Abstract The present invention relates to novel compounds which have hemoregulatory activities and can be used to stimulate hematopoiesis and for the treatment of viral, fungal and bacterial infectious diseases.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

HEMOREGULATORY COMPOUNDS

Field of the Invention

5 The present invention relates to novel compounds which have hemoregulatory activities and can be used to stimulate hematopoiesis and for the treatment of viral, fungal and bacterial infectious diseases.

Background of the Invention

10 The hematopoietic system is a life-long cell renewal process whereby a defined stem cell population gives rise to a larger population of mature, differentiated blood cells (Dexter TM. Stem cells in normal growth and disease. Br Med J 1987; 195:1192-1194) of at least nine different cell lineages (erythrocytes, platelets, eosinophils, basophils, neutrophils,
15 monocytes/macrophages, osteoclasts, and lymphocytes) (Metcalf D. The Molecular Control of Blood Cells. 1988; Harvard University Press, Cambridge, MA). Stem cells are also ultimately responsible for regenerating bone marrow following treatment with cytotoxic agents or following bone marrow transplantation.

20 The major dose-limiting toxicities of most standard anti-neoplastic drugs are related to bone marrow suppression, which if severe and prolonged, can give rise to life-threatening infectious and hemorrhagic complications. Myelosuppression is predictable and has been reported to be dose-limiting in
25 greater than 50% of single-agent Phase I trials cytotoxic compounds (Merrouche Y, Catimel G, Clavel M. Hematopoietic growth factors and chemoprotectants; should we move toward a two-step process for phase I clinical trials in oncology? Ann Oncol 1993; 4:471-474). The risk of infection is directly related to the degree of myelosuppression as measured by the
30 severity and duration of neutropenia (Brody GP, Buckley M, Sathe YS, Freireich EJ. Quantitative relationship between circulating leukocytes and infections with acute leukemia. Ann In Med 1965; 64:328-334).

The control of hematopoiesis involves the interplay of a variety of cytokines and growth factors during various stages of the hematopoietic cascade, including early pluripotent stem cells and mature circulating effector
5 cells. These regulatory molecules include granulocyte colony stimulating factor (G-CSF), granulocyte-macrophage stimulating factor (GM-CSF), macrophage-colony stimulating factor (M-CSF), and a variety of interleukins which have overlapping, additive and synergistic actions which play major roles in host defence. Mechanistically, this is accomplished by enhancing the
10 production of granulocytes and macrophages, as well as by the activation of effector cell functions (Moore MAS. Hemopoietic growth factor interactions: in vitro and in vivo preclinical evaluation. Cancer Surveys 1990; 9:7-80). These coordinated activities support optimal host defences which are necessary for fighting bacterial, viral and fungal infections.

15
Strategies to prevent or reduce the severity of neutropenia and myelotoxicity include the use of hematopoietic growth factors and/or other hematopoietic cytokines. Such treatments are becoming common practice, in that they offer the potential of increased doses of cytotoxic agents that may
20 improve the therapeutic efficacy of antineoplastic agents, and reduce the morbidity associated with their use (Steward WP. Granulocyte and granulocyte-macrophage colony stimulating factors, Lancet 1993; 342:153-157). Clinical studies have demonstrated the G-, GM- and/or M-CSF may reduce the duration of neutropenia, accelerate myeloid recovery, and reduce
25 neutropenia-associated infections and other infectious complications in patients with malignancies who are receiving cytotoxic chemotherapy or in high infectious-risk patients following bone marrow transplantation (Steward WP. Granulocyte and granulocyte-macrophage colony stimulating factors, Lancet 1993; 342:153-157 and Munn DH, Cheung NKV. Preclinical and
30 clinical studies of macrophage colony-stimulating factor. Semin Oncol 1992; 19:395-407).

Synthetic peptides have been reported to induce the synthesis and release of haematopoietic mediators, including m-CSF from bone marrow stromal elements see U.S. Patent Application 08/001,905.

5 We have now found certain novel non-peptide compounds which have a stimulative effect on myelopoietic cells. They are useful in stimulating myelopoiesis in patients suffering from reduced myelopoietic activity, including bone marrow damage, agranulocytosis and aplastic anemia including patients having depressed bone marrow function due to
10 immunosuppressive treatment to suppress tissue reactions i.e. in bone marrow transplant surgery. They may also be used to promote more rapid regeneration of bone marrow after cytostatic chemotherapy and radiation therapy for neoplastic and viral diseases. They may be of particular value where patients have serious infections due to a lack of immune response
15 following bone marrow failure. They are useful in the treatment and prevention of viral, fungal and bacterial disease.

Summary of the Invention

 This invention comprises compounds, hereinafter represented as
20 Formula (I), which have hemoregulatory activities and can be used to stimulate hematopoiesis and in the prevention and treatment of bacterial, viral and fungal diseases.

 These compounds are useful in the restoration of leukocytes in patients
25 with lowered cell counts resulting from a variety of clinical situations, such as surgical induced myelosuppression, AIDS, ARDS, congenital myelodysplasia, bone marrow and organ transplants; in the protection of patients with leukopenia from infection; in the treatment of severely burned patients and in the amelioration of the myelosuppression observed with some cell-cycle
30 specific antiviral agents and in the treatment of infections in patients who have had bone marrow transplants, especially those with graft versus host disease, in the treatment of tuberculosis and in the treatment of fevers of unknown

origin in humans and animals. The compounds are also useful in the treatment and prevention of viral, fungal and bacterial infectious diseases, particularly Candida and Herpes in both immunosuppressed and "normal" subjects. They are useful in the treatment of sepsis caused by gram negative
5 and gram positive organisms.


These compounds may also be used in combination with the myelosuppressive agents of co-pending U.S. Application No. 07/799,465 and U.S. Patent No. 4,499,081, incorporated by reference herein, to provide
10 alternating peaks of high and low activity in bone marrow cells, thus augmenting the natural circadian rhythm of hematopoiesis. In this way, cytostatic therapy can be given at periods of low bone marrow activity, thus reducing the risk of bone marrow damage, while regeneration will be promoted by the succeeding peak of activity. This invention is also a
15 pharmaceutical composition, which comprises a compound of Formula (I) and a pharmaceutically acceptable carrier.

This invention further constitutes a method for stimulating the myelopoietic system of an animal, including humans, which comprises
20 administering to an animal in need thereof, an effective amount of a compound of Formula (I).

This invention also constitutes a method for preventing and treating viral, fungal and bacterial infections in immunosuppressed and normal
25 animals, including humans, which comprises administering to an animal in need thereof, an effective amount of a compound of Formula (I).

Detailed Description of the Invention

The compounds of the invention are represented by structural Formula I



A^1 and A^2 are independently $\text{Z}-(\text{CH}_2)_k-(\text{NR}^2)_y-$.

Z is independently a 4 - 10 membered mono- or bicyclic heterocyclic ring system containing up to four heteroatoms N, O, S in the ring in which at least one heteroatom is N, and wherein the ring is substituted or unsubstituted by one or two C_{1-4} alkyl, F, Cl, Br, I, C_{1-4} alkoxy, $(\text{CH}_2)_m\text{R}_4$, oxo, oxime, $\text{O}-\text{C}_{1-4}$ alkyloxime, hydroxy, $\text{N}(\text{R}_3)_2$, acylamino or aminoacyl groups, 8, 9, 10 membered monocyclic ring systems being excluded;

R^1 and R^2 are independently hydrogen, C_{1-4} alkyl, $\text{C}(\text{O})\text{R}_4$, C_{1-4} alkyl or R_1 and R_2 are benzyl which is optionally substituted by one or two C_{1-4} alkyl, C_{1-4} alkoxy, F, Cl, I, Br, OH, or $\text{N}(\text{R}_3)_2$;

R_3 is independently hydrogen, C_{1-4} alkyl, or benzyl;

R_4 is independently OR_3 , $\text{N}(\text{R}_3)_2$ or SR_3 ; and

k is an integer from 0 to 4;

m is independently an integer from 1 to 3;

n is 1 or 2;

y is zero or one;

or a pharmaceutically acceptable salt thereof.

C_{1-4} alkyl groups may be straight or branched.

The compounds of the present invention may contain one or more asymmetric carbon atoms and may exist in racemic and optically active form. All these compounds and diastereomers are contemplated to be within the scope of the present invention.

Z in the above Formula (I) denotes an optionally substituted pyrrolyl, isopyrrolyl, pyrazolyl, isoimidazolyl, triazolyl, iosxazolyl, oxazolyl, thiazolyl, isothiazolyl, oxadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, pyrrolidinyl, piperazinyl, triazinyl, morpholinyl, indolyl, indoleninyl, isobenzazolyl, pyrindinyl, ioindazolyl, indoxazinyl, benzoxazolyl, quinolinyl, isoquinolinyl, cinnolinyl, quinazolinyl, naphthyridinyl, pyridopyridinyl, tetrahydroquinolinyl, tetrahydroisoquinolinyl, quinoxalinyl, indolinyl, 2-pyrrolidonyl, imidazolyl, imidazolidinyl, imidazoliny, piperidyl, tetrazolyl, quinuclidinyl, azetidiny, or purinyl;

10 Preferred compounds are those wherein Z is optionally substituted pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl, quinolinyl, tetrahydroquinolinyl, azetidiny, or pyrrolidinyl;


More preferred compounds are those wherein Z is optionally substituted 2-pyridinyl, 2-pyrimidinyl, 2-pyrazinyl, 2-pyrrolidon-5-yl, or pyrrolidinyl.

15 Preferred compounds are N,N'-bis(picolinoyl)-1,4-diazacycloheptane, N,N'-bis(picolinoyl)piperazine, N,N'-bis(pyroglutamoyl)piperazine or N,N'-bis(pyroglutamoyl)-1,4-diazacycloheptane.

20

Methods of Preparation

Compounds of Formula (I) wherein A¹, A² and m are defined as in Formula (I) are prepared by methods analogous to those described in Scheme 3. Appropriate diamines (such as **1** in Scheme 3) are bis-acylated with a suitable heterocyclic acid (such as **2** in Scheme 3) using an activating agent (such as BOP reagent) with a suitable base (such as iPr₂NEt) in a polar aprotic solvent (such as DMF) to give final product.

Scheme 1

a) BOP, HOBT, iPr, NEt, DMF

5

In order to use a compound of the Formula (I) or a pharmaceutically acceptable salt thereof for the treatment of humans and other mammals it is normally formulated in accordance with standard pharmaceutical practice as a pharmaceutical composition.

10

According to a still further feature of the present invention there are provided pharmaceutical compositions comprising as active ingredient one or more compounds of Formula (I) as herein before defined or physiologically compatible salts thereof, in association with a pharmaceutical carrier or excipient. The compositions according to the invention may be presented for example, in a form suitable for oral, nasal, parenteral or rectal administration.

15

As used herein, the term "pharmaceutical" includes veterinary applications of the invention. These compounds may be encapsulated, tableted or prepared in an emulsion or syrup for oral administration. Pharmaceutically acceptable solid or liquid carriers may be added to enhance or stabilize the composition, or to facilitate preparation of the composition. Liquid carriers include syrup, peanut oil, olive oil, glycerin, saline and water. Solid carriers include starch, lactose, calcium sulfate dihydrate, terra alba, magnesium stearate or stearic acid, talc, pectin, acacia, agar or gelatin. The carrier may also include a sustained release material such a glyceryl monostearate or glyceryl distearate, alone or with a wax. The amount of

20

25

solid carrier varies but, preferably will be between about 20 mg to about 1 g per dosage unit. The pharmaceutical preparations are made following the conventional techniques of pharmacy involving milling, mixing, granulating, and compressing, when necessary, for tablet forms; or milling, mixing and filling for hard gelatin capsule forms. Capsules containing one or several active ingredients may be produced, for example, by mixing the active ingredients with inert carriers, such as lactose or sorbitol, and filling the mixture into gelatin capsules. When a liquid carrier is used, the preparation will be in the form of a syrup, elixir, emulsion or an aqueous or non-aqueous suspension. Such a liquid formulation may be administered directly p.o. or filled into a soft gelatin capsule. Organ specific carrier systems may also be used.

Alternately pharmaceutical compositions of the compounds of this invention, or derivatives thereof, may be formulated as solutions of lyophilized powders for parenteral administration. Powders may be reconstituted by addition of a suitable diluent or other pharmaceutically acceptable carrier prior to use. The liquid formulation is generally a buffered, isotonic, aqueous solution. Examples of suitable diluents are normal isotonic saline solution, standard 5% dextrose in water or buffered sodium or ammonium acetate solution. Such formulation is especially suitable for parenteral administration, but may also be used for oral administration and contained in a metered dose inhaler or nebulizer for insufflation. It may be desirable to add excipients such as polyvinylpyrrolidone, gelatin, hydroxy cellulose, acacia, polyethylene glycol, mannitol, sodium chloride or sodium citrate.

For rectal administration, a pulverized powder of the compounds of this invention may be combined with excipients such as cocoa butter, glycerin, gelatin or polyethylene glycols and molded into a suppository. The pulverized powders may also be compounded with an oily preparation, gel, cream or

emulsion, buffered or unbuffered, and administered through a transdermal patch.

5 Nasal sprays may be formulated similarly in aqueous solution and packed into spray containers either with an aerosol propellant or provided with means for manual compression.

Dosage units containing the compounds of this invention preferably contain .05-50 mg, for example .05-5 mg of the compound of formula (I) or
10 salt thereof.

According to a still further feature of the present invention there is provided a method of stimulation of myelopoiesis which comprises administering an effective amount of a pharmaceutical composition as
15 hereinbefore defined to a subject.

No unacceptable toxicological effects are expected when compounds of the invention are administered in accordance with the present invention.

20 The biological activity of the compounds of Formula (I) are demonstrated by the following tests.

Induction of Hematopoietic Synergistic Activity in Stromal Cells

25 The murine bone marrow derived stromal cell line, C6.4 is grown in 12 well plates in RPMI 1640 with 10% FBS. Upon reaching confluence, the C6.4 cells are washed and the media exchanged with fresh RPMI 1640 without FBS. Confluent cell layers of murine C6.4 cells are treated with compound. Cell-free supernatants are collected 18 hours later. Supernatants
30 are fractionated with a Centricon-30 molecular weight cut-off membrane. C6.4 cell hematopoietic synergistic factor (HSF) activity is measured in a murine CFU-C assay.

CFU-C Assay

Bone marrow cells are obtained from C57B1/6 female mice and suspended in RPMI 1640 with 10% FBS. Bone marrow cells (7.5E+4
5 cells/mL) are cultured with sub optimal levels of CFU plus dilutions of test C6.4 cell 30K-E supernatants from above in a standard murine soft agar CFU-C assay. Cell aggregates >50 cells are counted as colonies. The number of agar colonies counted is proportional to the amount of HSF present within the C6.4 bone marrow stromal line supernatant.

10

Effector Cell Function Assay

Female C57B1 mice are administered test compound IP or PO daily for 8 days. Resident peritoneal exudate cells (PEC) utilized *ex vivo* from
15 treated or untreated mice are harvested with cold calcium and magnesium-free DPBS supplemented with heparin and antibiotics within 2-4 hours following the last injection. Adherent PEM populations are prepared by incubating standardized PEC suspensions in microtiter dishes for 2 hours at 37 °C (5% CO₂) and removing nonadherent cells by washing the wells with warm buffer.

20

The superoxide dismutase-inhibitable (SOD) superoxide released by effector cells in response to a *in vitro* stimulation by phorbol myristate acetate (PMA) (100-200nM) or pre-opsonized (autologous sera) live *C. albicans* (E:T = 1:10) are quantitated in a microtiter ferricytochrome *c* reduction assay. The
25 assay is performed in the presence of 1% gelatin/HBSS and 80uM ferricytochrome *c* in a total volume of 200uL/well. The nmoles of cytochrome *c* reduced/well is calculated from spectrophotometric readings (550 nm) taken following a 1 hour incubation at 37 °C (5% CO₂). The amount of SOD-inhibitable cytochrome *c* reduced is determined by the
30 inclusion of wells containing SOD (200 U/well). Baseline superoxide release

is determined in the absence of stimuli. Experimental data are expressed as a percentage of the control group.

The following examples are illustrative and are not limiting of the
5 compounds of this invention.

EXAMPLE 1

N,N'-Bis(picolinoyl)-1,4-diazacycloheptane

To a solution of 1,4-diazacycloheptane (0.31 g, 3.08 mmol) in pyridine
10 (30 mL) was added picolinic acid (0.74 g, 6.01 mmol) and EDC (1.16 g, 6.05 mmol). After 18 h at room temperature, the bulk of the pyridine was removed *in vacuo*. The residue was dissolved in water (50 mL) and extracted with EtOAc (3 x 50 mL). The combined organic extracts were washed with water (50 mL) and dried over Na₂SO₄. Removal of solvent gave 0.14 g of a red
15 syrup. Purification by flash chromatography (20 % MeOH/EtOAc, silica gel) afforded 0.09 g (10 %) of the title compound. MS (ES+) m/z 311.2 [M+H]⁺.

EXAMPLE 2

N,N'-Bis(picolinoyl)piperazine

20

In a manner analogous to Example 1, piperazine (0.09 g, 1.04 mmol), picolinic acid (0.37 g, 3.00 mmol) and EDC (0.57 g, 3.00 mmol) in pyridine (10 mL) gave 0.29 g (94 %) of the title compound. MS (ES+) m/z 297.0 [M+H]⁺.

25

EXAMPLE 3

Formulations for pharmaceutical use incorporating compounds of the present invention can be prepared in various forms and with numerous excipients. Examples of such formulations are given below.

<u>Tablets/Ingredients</u>	<u>Per Tablet</u>
1. Active ingredient (Cpd of Form.I)	0.5 mg
2. Corn Starch	20 mg
5 3. Alginic acid	20 mg
4. Sodium alginate	20 mg
5. Mg stearate	1.3 mg

Procedure for tablets:


- Step 1 Blend ingredients No. 1, No. 2, No. 3 and No. 4
 10 in a suitable mixer/blender.
- Step 2 Add sufficient water portion-wise to the blend
 from Step 1 with careful mixing after each
 addition. Such additions of water and mixing
 until the mass is of a consistency to permit
 15 its conversion to wet granules.
- Step 3 The wet mass is converted to granules by passing
 it through an oscillating granulator using a
 No. 8 mesh (2.38 mm) screen.
- Step 4 The wet granules are then dried in an oven at
 20 140°F (60°C) until dry.
- Step 5 The dry granules are lubricated with ingredient No. 5.
- Step 6 The lubricated granules are compressed on a suitable tablet press.

Parenteral Formulation

- 25 A pharmaceutical composition for parenteral administration is
 prepared by dissolving an appropriate amount of a compound of Formula I in
 polyethylene glycol with heating. This solution is then diluted with water for
 injections Ph Eur. (to 100 ml). The solution is then sterilized by filtration
 30 through a 0.22 micron membrane filter and sealed in sterile containers.

CLAIMS:

1. A compound of Formula (I)



A^1 and A^2 are independently $\text{Z}-(\text{CH}_2)_k-(\text{NR}^2)_y-$.

- 10 Z is independently a 4 - 10 membered mono- or bicyclic heterocyclic ring system containing up to four heteroatoms N, O, S in the ring in which at least one heteroatom is N, and wherein the ring is substituted or unsubstituted by one or two C_{1-4} alkyl, F, Cl, Br, I, C_{1-4} alkoxy, $(\text{CH}_2)_m\text{R}_4$, oxo, oxime, $\text{O}-\text{C}_{1-4}$ alkyloxime, hydroxy, $\text{N}(\text{R}_3)_2$, acylamino or aminoacyl groups, 8, 9, 10 membered monocyclic ring systems being excluded;

- 15 R^1 and R^2 are independently hydrogen, C_{1-4} alkyl $\text{C}(\text{O})\text{R}_4$, C_{1-4} alkyl or R^1 and R^2 are benzyl which is optionally substituted by one or two C_{1-4} alkyl, C_{1-4} alkoxy, F, Cl, I, Br, OH, or $\text{N}(\text{R}_3)_2$;

R_3 is independently hydrogen, C_{1-4} alkyl, or benzyl;

R_4 is independently OR_3 , $\text{N}(\text{R}_3)_2$ or SR_3 ; and

- 20 k is an integer from 0 to 4;

m is independently 2 or 3;

y is zero or one;

or a pharmaceutically acceptable salt thereof.

- 25 2. A compound of Claim 1 wherein Z is optionally substituted pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl, quinolinyl, tetrahydroquinolinyl, or pyrrolidinyl.

3. A compound of Claim 2 wherein Z is optionally substituted 2-pyridinyl, 2-pyrimidinyl, 2-pyrazinyl, 2-pyrrolidon-5-yl, or 2-pyrrolidinyl.

4. A compound of Claim 1 selected from the group consisting of N,N'-bis(picolinoyl)-1,4-diazacycloheptane, N,N'-bis(picolinoyl)piperazine, N,N'-bis(pyroglutamoyl)piperazine or N,N'-bis(pyroglutamoyl)-1,4-diazacycloheptane.

5. A pharmaceutical composition comprising a compound of Formula (I) and a pharmaceutical carrier or excipient.

6. A method of preventing or treating viral, fungal or bacterial infections which comprises administering to an animal in need thereof, an effective amount of a compound of Formula I.

7. A method of stimulating myelopoiesis in an animal in need thereof by administering a compound of claim 1.

8. A method of treating or preventing sepsis in an animal in need thereof by administering a compound of claim 1.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US96/18342

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) :A61K, 31/55, 31/495; C07D, 401/14, 403/14

US CL :Please See Extra Sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 540-375; 544, 359, 360, 363, 372

514, 219, 252

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Please See Extra Sheet.

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US, A 5,292,726 (ASHTON et al.) 08 March 1994 (08.03.94) (see the entire document)	1-8

☐ Further documents are listed in the continuation of Box C.
 ☐ See patent family annex.

* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document published on or after the international filing date "L" document which may throw doubt on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed		"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "Z" document member of the same patent family
--	--	--

Date of the actual completion of the international search

12 FEBRUARY 1997

Date of mailing of the international search report

05 MAR 1997

 Name and mailing address of the ISA/US
 Commissioner of Patents and Trademarks
 Box PCT
 Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

ROBERT T. BOND aco

Telephone No. (703) 308-1235

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US96/18342

A. CLASSIFICATION OF SUBJECT MATTER:

US CL :

540, 375; 544, 359, 360, 363, 372

514, 219, 252

B. FIELDS SEARCHED

Documentation other than minimum documentation that are included in the fields searched:

CHEMICAL ABSTRACTS 1-4-DIAZEPINE VOL. 32—> VOL.124
(1938 —> JUNE 1996)

HOMOPIPERAZINE VOL. 21 -> VOL. 31 (1427-1937)